




## Potato

<b>Culture and Variety</b>	<b>:</b>	<b>AICRP-P-9 (Kufri Ganga)</b>	
Status of release		Released by CVRC from Shimla for Odisha, with involvement of Scientistis of AICRP on Potato, OUAT	
Duration		75-80 days at Bhubaneswar	
Potential yield		32 t/ha	
Adaptability		All potato growing areas of Odisha	
Advantage/ Superiority over existing ruing Variety		45% higher yield over KufriJyoti (22 t/ha)	
Other characteristics		white skinned, table purpose, moderately heat tolerant, needssecondearthing up as tubers get exposed above soil	
<b>Culture and Variety</b>	<b>:</b>	<b>AICRP-C-28 (Kufri Lima)</b>	
Status of release		Released by CVRC from Shimla for Odisha, with involvement of Scientistis of AICRP on Potato, OUAT	
Duration		90 days at Bhubaneswar	
Potential yield		30 t/ha	
Adaptability		All potato growing areas of Odisha, where bacterial wilt is absent or mild.	
Advantage/ Superiority over existing ruing Variety		36% higher yield over KufriJyoti (22 t/ha)	
Other characteristics		Dark green glossy leaves, virus resistant, white skinned, heattolerant	
<b>Culture and Variety</b>	<b>:</b>	<b>AICRP-P-14 (KufriNeelkanth)</b>	
Status of release		Released by CVRC from Shimla for Odisha, with involvement of Scientistis of AICRP on Potato, OUAT	
Duration		75 days at Bhubaneswar	
Potential yield		28 t/ha	
Adaptability		All potato growing areas of Odisha	
Advantage/ Superiority over existing ruing Variety		27% higher yield over KufriJyoti (22 t/ha)	
Other characteristics		Purple skinned, table purpose, rich in carotenoids, anthocyaninand anti-oxidants	
Importance		More preferred for ill and weak people	